


Abstract

The present invention provides compounds of formula (I) including stereoisomers, prodrugs and pharmaceutically acceptable salts or solvates thereof



wherein

the dashed line may represent a double bond;

R is aryl or heteroaryl, each of which may be substituted by 1 to 4 groups J selected from:

halogen, C1-C6 alkyl, C1-C6 alkoxy, halo C1-C6 alkyl, C2-C6 alkenyl, C2-C6 alkynyl, halo C1-C6 alkoxy, -C(O)R₂, nitro, hydroxy, -NR₃R₄, cyano, and or a group Z;

R₁ is hydrogen, C3-C7 cycloalkyl, C1-C6 alkyl, C1-C6 alkoxy, C1-C6 thioalkyl, C2-C6 alkenyl, C2-C6 alkynyl, halo C1-C6 alkyl, halo C1-C6 alkoxy, halogen, NR₃R₄ or cyano;

R₂ is a C1-C4 alkyl, -OR₃ or -NR₃R₄;

R₃ is hydrogen or C1-C6 alkyl;

R₄ is hydrogen or C1-C6 alkyl;

R₅ is a C1-C6 alkyl, halo C1-C6 alkyl, C1-C6 alkoxy, halo C1-C6 alkoxy, C3-C7 cycloalkyl, hydroxy, halogen, nitro, cyano, -NR₃R₄; -C(O)R₂;

R₆ is a C1-C6 alkyl, halo C1-C6 alkyl, C1-C6 alkoxy, halo C1-C6 alkoxy, C3-C7 cycloalkyl, hydroxy, halogen, nitro, cyano, -NR₃R₄; -C(O)R₂;

R₇ is hydrogen, C1-C6 alkyl, halogen or halo C1-C6 alkyl;

R₈ is hydrogen, C3-C7 cycloalkyl, C1-C6 alkyl, C2-C6 alkenyl, C2-C6 alkynyl, NR₃R₄ or cyano;

R₉ is hydrogen, C3-C7 cycloalkyl, C1-C6 alkyl, C2-C6 alkenyl, C2-C6 alkynyl, NR₃R₄ or cyano;

R₁₀ is hydrogen, C3-C7 cycloalkyl, C1-C6 alkyl, C2-C6 alkenyl, C2-C6 alkynyl, NR₃R₄ or cyano;

R₁₁ is hydrogen, C3-C7 cycloalkyl, C1-C6 alkyl, C2-C6 alkenyl, C2-C6 alkynyl, NR₃R₄ or cyano;

R₁₂ is hydrogen, C3-C7 cycloalkyl, C1-C6 alkyl, C2-C6 alkenyl, C2-C6 alkynyl, NR₃R₄ or cyano;

R₁₃ is hydrogen, C3-C7 cycloalkyl, C1-C6 alkyl, C2-C6 alkenyl, C2-C6 alkynyl, NR₃R₄ or cyano;

R₁₄ is R₃ or -C(O)R₂;

D is CR₈R₉ or is CR₈ when double bonded with G or A;

- G is $\text{CR}_{10}\text{R}_{11}$ or is CR_{10} when double bonded with D or is CR_{10} when double bonded with X when X is carbon;
- A is $\text{CR}_{12}\text{R}_{13}$ or is CR_{12} when double bonded with D;
- X is carbon or nitrogen;
- Y is nitrogen or $-\text{CR}_7$;
- W is a 4-8 carbocyclic membered ring, which may be saturated or may contain one to three double bonds, and

in which:

- one carbon atom is replaced by a carbonyl or $\text{S}(\text{O})_m$; and
- one to four carbon atoms may optionally be replaced by oxygen, nitrogen or NR_{14} , $\text{S}(\text{O})_m$, carbonyl, and such ring may be further substituted by 1 to 8 R_6 groups;

- Z is a 5-6 membered heterocycle or a phenyl, which may be substituted by 1 to 8 R_5 groups;

- m is an integer from 0 to 2,

to processes for their preparation, to pharmaceutical compositions containing them and to their use in the treatment of conditions mediated by corticotropin-releasing factor (CRF).